

Everwilde Spring Dry Mix

Native Wildflowers

| | Latin Name | Common Name | Seeds/Oz | Bloom Time & Color | | | | | | | Height | Light | | | Soil | | | Percent By | | |
|----|--------------------------|---------------------------|----------|--------------------|---|---|---|---|---|---|--------|-------|---|---|------|---|---------------|---------------|--------|-------|
| | | | | A | M | J | J | A | S | O | | N | S | P | F | W | M | D | Weight | Seeds |
| 1 | Allium stellatum | Prairie Onion | 11,000 | | | J | J | A | | | | | | P | F | | M | D | 4.17% | 2.50% |
| 2 | Amorpha canescens | Lead Plant | 16,000 | | | J | J | A | | | | | P | F | | M | D | 2.50% | 2.18% | |
| 3 | Anemone cylindrica | Thimbleweed | 26,000 | M | J | | | | | | | | P | F | | M | D | 0.83% | 1.18% | |
| 4 | Artemisia ludoviciana | Prairie Sage | 250,000 | | | J | J | A | S | | | | P | F | | M | D | 0.17% | 2.28% | |
| 5 | Asclepias syriaca | Common Milkweed | 4,000 | | | J | J | A | | | | | P | F | | M | D | 3.33% | 0.73% | |
| 6 | Asclepias tuberosa | Butterfly Weed | 4,300 | | | J | J | A | | | | | P | F | | M | D | 1.25% | 0.29% | |
| 7 | Aster dumosus | Rice Button Aster | 57,000 | | | | | A | S | O | | | | F | | M | D | 0.83% | 2.59% | |
| 8 | Baptisia alba | White Wild Indigo | 1,700 | | | J | J | | | | | | P | F | | M | D | 2.50% | 0.23% | |
| 9 | Baptisia leucophaea | Cream Wild Indigo | 1,400 | M | J | | | | | | | | P | F | | M | D | 0.83% | 0.06% | |
| 10 | Baptisia tinctoria | Yellow Wild Indigo | 4,000 | | | J | J | A | S | | | | P | F | | M | D | 1.25% | 0.27% | |
| 11 | Campanula rotundifolia | Harebell | 900,000 | | | J | J | | | O | N | | P | F | | M | D | 0.08% | 4.10% | |
| 12 | Cassia fasciculata | Partridge Pea | 2,700 | | | J | J | A | S | | | | P | F | | M | D | 8.33% | 1.23% | |
| 13 | Coreopsis lanceolata | Sand Coreopsis | 20,000 | | | J | J | A | | | | | | F | | | D | 2.50% | 2.73% | |
| 14 | Desmodium sessilifolium | Sessile-leaf Tick Trefoil | 5,000 | | | J | J | A | S | | | | P | F | | M | D | 0.63% | 0.17% | |
| 15 | Echinacea purpurea | Purple Coneflower | 6,600 | | | J | J | A | S | | | | P | F | | M | D | 6.67% | 2.40% | |
| 16 | Echinacea tenesseeensis | Tennessee Coneflower | 6,000 | | | J | J | A | S | | | | P | F | | M | D | 1.67% | 0.55% | |
| 17 | Helianthus giganteus | Tall Sunflower | 10,000 | | | J | J | A | S | | | | P | F | | M | D | 0.83% | 0.46% | |
| 18 | Heuchera richardsonii | Prairie Alumroot | 700,000 | M | J | J | | | | | | | P | F | | M | D | 0.08% | 3.19% | |
| 19 | Lespedeza capitata | Round-Headed Bush Clover | 8,000 | | | | | A | S | | | | P | F | | M | D | 3.33% | 1.46% | |
| 20 | Liatris aspera | Button Blazing Star | 16,000 | | | J | J | A | S | O | | | P | F | | M | D | 3.33% | 2.91% | |
| 21 | Lupinus perennis | Wild Lupine | 1,100 | M | J | | | | | | | | P | F | | | D | 1.67% | 0.10% | |
| 22 | Monarda punctata | Spotted Bee Balm | 90,000 | | | J | J | A | | | | | P | F | | | D | 0.63% | 3.07% | |
| 23 | Oenothera biennis | Common Evening Primrose | 90,000 | | | J | J | A | S | | | | P | F | | M | D | 0.63% | 3.07% | |
| 24 | Psoralea onobrychis | French Grass | 1,200 | | | J | J | A | | | | | | F | | | D | 3.33% | 0.22% | |
| 25 | Ratibida columnifera | Long-Headed Coneflower | 42,000 | | | J | J | A | | | | | P | F | | M | D | 1.25% | 2.87% | |
| 26 | Ratibida pinnata | Yellow Coneflower | 30,000 | | | J | J | A | S | | | | P | F | | M | D | 1.67% | 2.73% | |
| 27 | Rosa carolina | Pasture Rose | 2,500 | | | J | J | A | | | | | P | F | | M | D | 2.50% | 0.34% | |
| 28 | Rudbeckia hirta | Black-Eyed Susan | 92,000 | | | J | J | A | S | O | | | P | F | | M | D | 0.63% | 3.14% | |
| 29 | Tradescantia ohioensis | Ohio Spiderwort | 8,000 | M | J | J | | | | | | | P | F | | M | D | 0.83% | 0.36% | |
| 30 | Veronicastrum virginicum | Culver's Root | 800,000 | | | J | J | A | | | | | P | F | | M | D | 0.08% | 3.64% | |
| | | | | | | | | | | | | | | | | | 58.33% | 51.07% | | |

Native Grasses & Sedges

| | Latin Name | Common Name | Seeds/Oz | Bloom Time & Color | | | | | | | Height | Light | | | Soil | | | Percent By | |
|---|------------------------|---------------------|----------|--------------------|---|---|---|---|---|---|--------|-------|---|---|------|---|---------------|---------------|--------|
| | | | | A | M | J | J | A | S | O | | N | S | P | F | W | M | D | Weight |
| 1 | Andropogon scoparius | Little Bluestem | 15,000 | | | | | A | S | | | | P | F | | M | D | 12.50% | 10.24% |
| 2 | Bouteloua curtipendula | Side-Oats Grama | 6,000 | | | | | J | A | | | | P | F | | M | D | 4.17% | 1.37% |
| 3 | Carex brevoir | Plains Oval Sedge | 29,000 | | | J | J | | | | | S | P | F | | M | D | 4.17% | 6.60% |
| 4 | Muhlenbergia racemosa | Upland Wild Timothy | 80,000 | | | | | A | S | | | S | P | F | | | D | 2.08% | 9.10% |
| 5 | Panicum virgatum | Switch Grass | 14,000 | | | | | A | S | | | | P | F | | M | D | 10.42% | 7.97% |
| 6 | Sporobolus asper | Rough Dropseed | 30,000 | | | J | J | | | | | | P | F | | M | D | 8.33% | 13.65% |
| | | | | | | | | | | | | | | | | | 41.67% | 48.93% | |